(Biomedical Applications, Vol. 87)

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, ASCA, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Clinical Chemistry Lookout, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Pharmaceutical Abstracts, Referativnyi Zhurnal, Science Citation Index and Trends in Biotechnology)

Quantitative profiling of 6-ketoprostaglandin $F_{1\alpha}$, 2,3-dinor-6-ketoprostaglandin $F_{1\alpha}$, thromboxane B_2 and 2,3-dinor-thromboxane B_2 in human and rat urine by immunoaffinity extraction with gas chromatography–mass spectrometry
by C. Chiabrando, V. Pinciroli and A. Campoleoni (Milan, Italy), A. Benigni and A. Piccinelli (Bergamo, Italy) and R. Fanelli (Milan, Italy) (Received May 29th, 1989) 1

Specific and sensitive quantitation of medroxyprogesterone acetate in human serum by gas chromatography–mass spectrometry
by T. Järvinen, T. Keinonen, S. Auriola and P. Peura (Kuopio, Finland) and E. Hirvonen and E. Palva (Helsinki, Finland) (Received June 7th, 1989) 13

Enantioselective measurement of the *Candida* metabolite D-arabinitol in human serum using multi-dimensional gas chromatography and a new chiral phase
by B. Wong and M. Castellanos (Cincinnati, OH, U.S.A.) (Received June 13th, 1989) 21

Carrier gas flow-rate and injection system in capillary gas chromatography of unconjugated and glycine-conjugated bile acids
by J.M. Street (Cincinnati, OH, U.S.A.) and D.J.H. Trafford and H.L.J. Makin (London, U.K.) (Received June 14th, 1989) 31

Determination of leucine enkephalin and methionine enkephalin in equine cerebrospinal fluid by microbore high-performance liquid chromatography and capillary zone electrophoresis coupled to tandem mass spectrometry
by W.M. Mück and J.D Henion (Ithaca, NY, U.S.A.) (Received May 27th, 1989) .. 41

High-performance liquid chromatographic determination of some polar phospholipids in serum
by J. Böswart (Prague, Czechoslovakia), T. Schmidt (Leipzig, G.D.R.) and P. Kostiuk, V. Pacáková and K. Štulik (Prague, Czechoslovakia) (Received June 22nd, 1989) 61

Separation of tryptophan metabolites by reversed-phase high-performance liquid chromatography with amperometric and fluorescence detection
by A.J. Elderfield, R.J.W. Truscott, I.E.T. Gan and G.M. Schier (Wollongong, Australia) (Received June 13th, 1989) 71

(Continued overleaf)

Contents (continued)

Determination of leucine and α-ketoisocaproic acid concentrations and specific activity in plasma and leucine specific activities in proteins using high-performance liquid chromatography
by F.F. Horber, J. Kahl, L. Lecavalier, B. Krom and M.W. Haymond (Rochester, MN, U.S.A.) (Received June 8th, 1989) 81

Determination of hippuric acid and furanic acid in serum of dialysis patients and control persons by high-performance liquid chromatography
by A. Pickert, A. Bäuerle and H.M. Liebich (Tübingen, F.R.G.) (Received June 6th, 1989) 95

Determination of serum cytidine deaminase activity using ion-pair reversed-phase liquid chromatography
by I.T. James, K. Herbert, D. Perrett and P.W. Thomson (London, U.K.) (Received June 8th, 1989) 105

Cell surface interactions with metal chelates
by H.G. Botros and M. Vijayalakshmi (Compiègne, France) (Received June 12th, 1989) 113

Gas chromatographic–mass spectrometric analysis of plasma nifedipine
by K.S. Patrick, E.J. Jarvi, A.B. Straughn and M.C. Meyer (Memphis, TN, U.S.A.) (Received June 8th, 1989) 123

Analysis of 4-hydroxycyclophosphamide by gas chromatography–mass spectrometry in plasma
by P.S. Hong and K.K. Chan (Los Angeles, CA, U.S.A.) (Received June 12th, 1989) 131

Automated reversed-phase chromatographic analysis of etoposide and teniposide in plasma by using on-line surfactant-mediated sample clean-up and column-switching
by M.A.J. van Opstal and F.A.L. van der Horst (Utrecht, The Netherlands), J.J.M. Holthuis (Amsterdam, The Netherlands) and W.P. van Bennedom and A. Bult (Utrecht, The Netherlands) (Received June 8th, 1989) 139

Determination of cocaine and selected metabolites in canine and human serum by reversed-phase high-performance liquid chromatography on coupled cyanopropyl and silica columns
by B.M. Lampert and J.T. Stewart (Athens, GA, U.S.A.) (Received June 16th, 1989) 153

Determination of artelinic acid in blood plasma by high-performance liquid chromatography
by O.R. Idowu, S.A. Ward and G. Edwards (Liverpool, U.K.) (Received June 12th, 1989) 167

Rapid high-performance liquid chromatographic methods for the determination of overdose concentrations of some non-steroidal anti-inflammatory drugs in plasma or serum
by P.J. Streete (London, U.K.) (Received May 26th, 1989) 179

High-performance liquid chromatographic determination of pirprofen and five of its metabolites in human plasma without hydrolysis and in human urine before and after chemical hydrolysis
by A. Sioufi, F. Marfil, J. Richard, D. Colussi and J.P. Dubois (Rueil-Malmaison, France) (Received June 2nd, 1989) 195

Determination of the enantiomeric composition of salsolinol in biological samples by high-performance liquid chromatography with electrochemical detection
by E. Pianezzola, V. Bellotti, E. Fontana and E. Moro (Milan, Italy) and J. Gal and D.M. Desai (Denver, CO, U.S.A) (Received May 29th, 1989) 205

Determination of pentoxifylline and its 5-hydroxy metabolite in human plasma by solid-phase extraction and high-performance liquid chromatography with ultraviolet detection
by G. Musch, T. Hamoir and D.L. Massart (Brussels, Belgium) (Received June 6th, 1989) 215

Determination of bromperidol in serum by automated column-switching high-performance liquid chromatography
by K. Hikida, Y. Inoue, T. Miyazaki, N. Kojima and Y. Ohkura (Fukuoka, Japan) (Received June 20th, 1989) 227

Notes

High-performance liquid chromatographic determination of malonaldehyde in serum
by S. Kawai and K. Kasashima (Gifu, Japan) and M. Tomita (Kurashiki, Japan) (Received May 27th, 1989) 235

Simultaneous determination of citrate and D-isocitrate in urine by isocratic ion chromatography
by R.P. Singh, S.A. Smesko and G.H. Nancollas (Buffalo, NY, U.S.A.) (Received June 13th, 1989) 239

Simple method for the determination of L-5-hydroxytryptophan in plasma by high-performance liquid chromatography
by G. Coppi and M. Barchielli (Milan, Italy) (Received June 13th, 1989) 245

Method for the determination of the specific activities of UTP and CTP in mouse kidney by high-performance liquid chromatography
by S.S. Elliger and G. Watson (Berkeley, CA, U.S.A.) (Received June 14th, 1989) . 249

Convenient method for the analysis of nicotinic acid as a metabolite of nicotinate esters in various tissue homogenates
by A. Durrer, B. Walther, A. Racciatti and B. Testa (Lausanne, Switzerland) (Received June 12th, 1989) 256

Simultaneous automated high-performance liquid chromatographic determination of 5-hydroxy-3-indoleacetic acid and homovanillic acid in urine with fluorescence detection
by M. Koel and P. Nebinger (Osnabrück, F.R.G.) (Received June 6th, 1989) 263

Improved high-performance liquid chromatographic technique for the determination of hepatic α-tocopherol
by H.R. Patel, S.P. Ashmore, L. Barrow and M.S. Tanner (Leicester, U.K.) (Received June 13th, 1989) 269

Determination of ximoprofen in human plasma by gas chromatography
by I.W. Taylor and L.F. Chasseaud (Huntingdon, U.K.) (Received June 2nd, 1989) 275

Gas chromatographic assay for chlorambucil and phenylacetic mustard in plasma
by F. Lohmar, E. Musch, G.E. Von Unruh and B.J. Mühlenbruch (Bonn, F.R.G.) (Received June 8th, 1989) 281

(Continued overleaf)

Contents (continued)

Determination of total 2,5-hexanedione by reversed-phase high-performance liquid chromatography
by I. Marchiseppe, M. Valentino and M. Governa (Ancona, Italy) and V. Stocchi (Urbino, Italy) (Received May 30th, 1989) 288

Determination of fluvoxamine in human plasma by high-performance liquid chromatography with ultraviolet detection
by J.P. Foglia, L.A. Birder and J.M. Perel (Pittsburgh, PA, U.S.A.) (Received June 2nd, 1989) 295

Determination of diclofenac in plasma and synovial fluid by high-performance liquid chromatography with electrochemical detection
by L. Zecca and P. Ferrario (Milan, Italy) (Received June 13th, 1989) 303

High-performance liquid chromatographic determination of indomethacin in human plasma and urine
by A. Avgerinos and S. Malamataris (Thessaloniki, Greece) (Received May 30th, 1989) 309

Determination of flumequine in biological fluids and meat by high-performance liquid chromatography
by L. Ellerbroek (Hannover, F.R.G.) and M. Bruhn (Göttingen, F.R.G.) (Received May 30th, 1989) 314

High-performance liquid chromatographic analysis of N,N′,N′′-triethylenethiophosphoramide in human plasma
by P.W. Tinsley, P.J. O'Dwyer and F.P. LaCreta (Philadelphia, PA, U.S.A.) (Received June 13th, 1989) 318

Automated high-performance liquid chromatographic assay for the β-carboline derivative Abecarnil in plasma
by W. Krause and G. Gröhler (Berlin, F.R.G.) (Received June 7th, 1989) 324

High-performance liquid chromatographic determination of mitomycin C in rat and human plasma and urine
by J.T. Dalton, E.R. Geuns and J.L.-S. Au (Columbus, OH, U.S.A.) (Received June 21st, 1989) 330

Isotachophoretic determination of sulbactam in rat serum
by I. Jelínek, V. Rejholec, V. Holá, Z. Roubal, J. Snopek and E. Smolková-Keulemansová (Prague, Czechoslovakia) (Received June 7th, 1989) 338

Determination of glycyrrhizin and its metabolite glycyrrhetinic acid in rabbit plasma by high-performance liquid chromatography after oral administration of licorzin
by X.Y. Zhang (Urumchi, China), R.J. Wu (Nanjing, China), J. Chen (Urumchi, China) and D.K. An (Nanjing, China) (Received May 30th, 1989) 343

Letters to the Editor

Determination of 5-hydroxypropafenone in biological fluids by fused-silica capillary gas chromatography using electron-capture detection
by G.L.-Y. Chan, J.E. Axelson, F.S. Abbott, K.M. McErlane and C.R. Kerr (Vancouver, Canada) (Received June 2nd, 1989) 349

Determination of ofloxacin in bronchoalveolar lavage fluid by high-performance liquid chromatography and fluorimetric detection
by K. Matsubayashi, T. Une and Y. Osada (Tokyo, Japan) (Received May 30th, 1989) 354

Automated high-performance liquid chromatography with column switching for on-line clean-up and analysis of diltiazem and metabolites in human plasma
by V. Ascalone and L. Flaminio (Limito, Italy) (Received June 14th, 1989) 358

Dextromethorphan phenotypes determined by high-performance liquid chromatography and fluorescence detection
by E. Jacqz-Aigrain, Y. Menard, M. Popon and H. Mathieu (Paris, France) (Received May 30th, 1989) 361

Related articles published in Journal of Chromatography, Vol. 479 364

Author Index 366

Subject index 370

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)